Meniscus Transplant Post-Operative Protocol

Phase I – Maximum Protection

Weeks 0 to 6:

- Brace 0-90 degrees for 6 weeks
 - o Recommend locking brace in extension for sleep
- Non-weight bearing with two crutches for 6 weeks
- Limit knee flexion to 90 degrees for 3 weeks; allow 90-120 degrees between 3 to 6 weeks

Goals

- o Reduce inflammation and pain
- o 0 degrees of knee extension

Exercise progression

- Emphasize patellofemoral mobilizations
- o Passive/active knee range of motion with 90° flexion limit
- o Quadriceps setting emphasize VMO function
- o Multi-plane straight leg raising
- o Open chain hip strengthening
- o Gait training

Phase II - Progressive Stretching and Early Strengthening

Weeks 6 to 8:

- Brace open to 0-90 degrees for ambulation
- Continue using two crutches, with gradual progression of weight bearing
 - o Increase WB to FWB over next 2 weeks

Exercise progression

- o Gradual progression to full knee flexion
- Continue to emphasize patella mobility
- o Begin bilateral closed kinetic chain strengthening limited range initially
- o Step-up progression
- Begin stationary bike with light resistance initially
- o Gait training normalize gait pattern

Phase III – Strengthening and Proprioceptive

Weeks 8 to 12:

Goals

o Full knee range of motion

Exercise progression

- Avoid rotational movements for 14 weeks
- Advance stationary biking program (increase intensity)
- o Introduce treadmill walking and elliptical trainer
- o Begin unilateral closed kinetic chain program
- Gym strengthening progression (leg press above 90-degrees, hamstrings curls etc.)

Phase IV- Advanced Strengthening and Plyometric Drills

Weeks 12 to 16:

Follow-up examination with the physician

<u>Goals</u>

 Pass return-to-sport functional test at > 90% (involved vs uninvolved limb) by 16 weeks

Exercise progression

- o Plyometric drills from bilateral to unilateral at 12 weeks
- o Linear running progression at 12 weeks
- o Progress to lateral and rotational stresses at 14 weeks
- o Multi-directional drills at 14 to 16 weeks

o Rev. 10/2017